CONTENTS

Special Issue: Forty Years of Computational Quantum Chemistry

Dedication viii
My life as a physicist: memories and perspectives
C.C.J. Roothaan (Chicago, IL, U.S.A.) 1
Computational quantum chemistry – then and now
G.G. Hall (Nottingham, U.K.) 13
Computational developments in SCF calculations
R.A. Poirier and C.D. Keefe (St John's, Nfld., Canada) 19
Configuration interaction method: the past and future perspectives
W. Duch (Toruń, Poland) 27
Effective Hamiltonian method for molecules
S. Huzinaga (Edmonton, Alta., Canada) 51
The prediction and calculation of properties of atoms in molecules
R.F.W. Bader and K.E. Laidig (Hamilton, Ont., Canada) 75
Determination and use of response densities
E. Kraka, J. Gauss and D. Cremer (Göteborg, Sweden) 95
Coupled Hartree–Fock calculations of magnetic properties of the benzene molecule: estimate of the Hartree–Fock limit for magnetic susceptibilities and nuclear shieldings
P. Lazzeretti, M. Malagoli and R. Zanasi (Modena, Italy) 127
Medium-size polarized basis sets for high-level-correlated calculations of molecular electric properties. III. Alkali (Li, Na, K, Rb) and alkaline-earth (Be, Mg, Ca, Sr) atoms
A.J. Sadlej and M. Urban (Lund, Sweden) 147
Finite field ab initio calculations of the dipole polarizabilities and first and second hyperpolarizabilities using multiple sets of polarization functions. I. The water molecule
J.E. Bloor (Knoxville, TN, U.S.A.) 173
Geometry and electronic structure studies using computational quantum chemistry
H. Basch (Ramat Gan, Israel) 185
Is the Mills–Nixon effect operative in perfluoro-benzo [1,2:3,4:5,6]tricyclobutene?
Z.B. Maksić, M. Eckert-Maksić and D. Kovaček (Zagreb, Croatia, Yugoslavia), M. Hodošček, K. Poljanec (Ljubljana, Slovenia, Yugoslavia) and J. Kudnig (Hamburg, F.R.G.) ... 201
On the structure and thermochemistry of [18]annulene
J.M. Schulman and R.L. Disch (Flushing, NY, U.S.A.) 213
A computational molecular orbital study of the oxide and oxepin valence tautomers of naphthalene
C.W. Bock, P. George and J.P. Glusker (Philadelphia, PA, U.S.A.) 227
The conjunction of aromaticity and hypervalency: the $C_5H_5SF_3$ molecule 1,1,1-trifluorothiabenzene[1]
Y. Xie, H.F. Schaefer III (Athens, GA, U.S.A.) and J.S. Thrasher (Tuscaloosa, AL, U.S.A.) 247
Ab initio studies of bonding trends. Part 8. The 26-electron A=B–C=D″ and the 30-electron A=B=C=D″ systems[1]
P. Pyykkö and N. Runeberg (Helsinki, Finland) 269
Ab initio studies of bonding trends. Part 9. The dicyanamide–carbon suboxide–dicyanoether–cyanogen azide isoelectronic series A=B=C=D=E
P. Pyykkö and N. Runeberg (Helsinki, Finland) 279
Ab initio molecular orbital study of 1-methylsilatrane and model compounds
A. Greenberg (New Brunswick, NJ, U.S.A.), C. Plant and C.A. Venanzi (Newark, NJ, U.S.A.) 291
The influence of *gem*-substitution on the stability of open-shell and closed-shell systems. The captodative and anomeric effects re-examined
G. Leroy, M. Sana and C. Wilante (Louvain-la-Neuve, Belgium) 303

CONTENTS

Special Issue: Forty Years of Computational Quantum Chemistry

Dedication viii
My life as a physicist: memories and perspectives
C.C.J. Roothaan (Chicago, IL, U.S.A.) 1
Computational quantum chemistry – then and now
G.G. Hall (Nottingham, U.K.) 13
Computational developments in SCF calculations
R.A. Poirier and C.D. Keefe (St John's, Nfld., Canada) 19
Configuration interaction method: the past and future perspectives
W. Duch (Toruń, Poland) 27
Effective Hamiltonian method for molecules
S. Huzinaga (Edmonton, Alta., Canada) 51
The prediction and calculation of properties of atoms in molecules
R.F.W. Bader and K.E. Laidig (Hamilton, Ont., Canada) 75
Determination and use of response densities
E. Kraka, J. Gauss and D. Cremer (Göteborg, Sweden) 95
Coupled Hartree–Fock calculations of magnetic properties of the benzene molecule: estimate of the Hartree–Fock limit for magnetic susceptibilities and nuclear shieldings
P. Lazzeretti, M. Malagoli and R. Zanasi (Modena, Italy) 127
Medium-size polarized basis sets for high-level-correlated calculations of molecular electric properties. III. Alkali (Li, Na, K, Rb) and alkaline-earth (Be, Mg, Ca, Sr) atoms
A.J. Sadlej and M. Urban (Lund, Sweden) 147
Finite field ab initio calculations of the dipole polarizabilities and first and second hyperpolarizabilities using multiple sets of polarization functions. I. The water molecule
J.E. Bloor (Knoxville, TN, U.S.A.) 173
Geometry and electronic structure studies using computational quantum chemistry
H. Basch (Ramat Gan, Israel) 185
Is the Mills–Nixon effect operative in perfluoro-benzo [1,2:3,4:5,6]tricyclobutene?
Z.B. Maksić, M. Eckert-Maksić and D. Kovaček (Zagreb, Croatia, Yugoslavia), M. Hodošček, K. Poljanec (Ljubljana, Slovenia, Yugoslavia) and J. Kudnig (Hamburg, F.R.G.) ... 201
On the structure and thermochemistry of [18]annulene
J.M. Schulman and R.L. Disch (Flushing, NY, U.S.A.) 213
A computational molecular orbital study of the oxide and oxepin valence tautomers of naphthalene
C.W. Bock, P. George and J.P. Glusker (Philadelphia, PA, U.S.A.) 227
The conjunction of aromaticity and hypervalency: the $C_5H_5SF_3$ molecule 1,1,1-trifluorothiabenzene[1]
Y. Xie, H.F. Schaefer III (Athens, GA, U.S.A.) and J.S. Thrasher (Tuscaloosa, AL, U.S.A.) 247
Ab initio studies of bonding trends. Part 8. The 26-electron A=B–C=D″ and the 30-electron A=B=C=D″ systems[1]
P. Pyykkö and N. Runeberg (Helsinki, Finland) 269
Ab initio studies of bonding trends. Part 9. The dicyanamide–carbon suboxide–dicyanoether–cyanogen azide isoelectronic series A=B=C=D=E
P. Pyykkö and N. Runeberg (Helsinki, Finland) 279
Ab initio molecular orbital study of 1-methylsilatrane and model compounds
A. Greenberg (New Brunswick, NJ, U.S.A.), C. Plant and C.A. Venanzi (Newark, NJ, U.S.A.) 291
The influence of *gem*-substitution on the stability of open-shell and closed-shell systems. The captodative and anomeric effects re-examined
G. Leroy, M. Sana and C. Wilante (Louvain-la-Neuve, Belgium) 303

Quantum-chemical study of the Fukui function as a reactivity index. Part 2. Electrophilic substitution on mono-substituted benzenes
W. Langenaeker, K. Demel and P. Geerlings (Brussels, Belgium) 329
The relative Lewis acidities of BF_3 and SO_3: the enthalpies of complexation with the anionic bases F^-, OH^-, NH_2^-, CH_3^- and CF_3^-
C.A. Deakyne (Charleston, IL, U.S.A.) and J.F. Liebman (Baltimore, MD, U.S.A.) 343
An ab initio molecular orbital study of the structure, energetics and bond activation of Al^+ complexes
M. Alcamí, O. Mó and M. Yáñez (Madrid, Spain) 357
Correlated proton transfer potentials. $(HO–H–OH)^-$ and $(H_2O–H–OH_2)^+$
Z. Latajka and S. Scheiner (Carbondale, IL, U.S.A.) 373
Efficient techniques for the decomposition of intermolecular interaction energy at SCF level and beyond
W.A. Sokalski and S.M. Roszak (Wroclaw, Poland) 387
Theoretical chemistry in solution. Some results and perspectives of the continuum methods and in particular of the polarizable continuum model
J. Tomasi, R. Bonaccorsi (Pisa, Italy), R. Cammi (Parma, Italy) and F.J. Olivares del Valle (Badajoz, Spain) 401
Theory of solvent effects on electronic spectra
H. Ågren (Uppsala, Sweden) and K.V. Mikkelsen (Copenhagen, Denmark) 425
Potential energy curves of the 5s Rydberg excited states and ionized states of Kr_2 studied by the SAC–CI theory
Y. Mizukami and H. Nakatsuji (Kyoto, Japan) 469
The $(H_2O)_2^*$ cluster at a geometry of intramolecular charge transfer
C.A. Nicolaides and P. Valtazanos (Athens, Greece) 483
A theoretical study of the neutral, protonated, and deprotonated trimers of HF and HCl
J.E. Del Bene (Youngstown, OH, U.S.A.) and I. Shavitt (Columbus, OH, U.S.A.) 499
Electronically excited states of C_3 in the CASSCF approach
J. Kalcher and R. Janoschek (Graz, Austria) 509
Ab initio CI calculation of the electronic states of HOOH: oscillator strengths and optical rotatory strengths
J. Hutter, H.P. Lüthi and T.-K. Ha (Zurich, Switzerland) 515
On quantum Monte Carlo for the electronic structure of molecules
B.L. Hammond, M.M. Soto, R.N. Barnett and W.A. Lester, Jr. (Berkeley, CA, U.S.A.) 525

Author index 539

Subject index 541